

¹ **special: A Python package for the spectral characterization of directly imaged low-mass companions**

³ **Valentin Christiaens**  ¹

⁴ 1 Space sciences, Technologies & Astrophysics Research Institute, Université de Liège, Belgium

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

5 **Summary**

6 Recent technological progress in high-contrast imaging has allowed the spectral characterization
7 of directly imaged giant planet and brown dwarf companions at ever shorter angular separation
8 from their host stars, hence opening a new avenue to study their formation, evolution, and
9 composition. In this context, **special** is a Python package that was developed to provide the
10 tools to analyse the low- to medium-resolution optical/IR spectra of these directly imaged
11 low-mass companions.

12 **Statement of need**

13 **special** provides the following tools for the analysis of measured spectra:

- 14 ▪ calculation of the spectral correlation between channels of an integral field spectrograph
15 (IFS) datacube ([Delorme et al., 2017](#); [Greco & Brandt, 2016](#));
- 16 ▪ calculation of empirical spectral indices for MLT-dwarfs ([Allers et al., 2007](#); [Gorlova et
al., 2003](#); [Slesnick et al., 2004](#)), enabling their classification;
- 17 ▪ fitting of input spectra to different (user-provided) grids of models, with the possibility
18 to include additional parameters such as extra blackbody component(s) and extinction;
- 19 ▪ estimating most likely model parameters in a Bayesian framework, using either MCMC
20 ([Goodman & Weare, 2010](#)) or nested ([Buchner, 2021a](#); [Feroz et al., 2009](#); [Mukherjee et
al., 2006](#); [Skilling, 2004](#)) samplers to infer their posterior distributions;
- 21 ▪ searching for the best-fit template spectrum within a given template library, with up to
22 two free parameters (flux scaling and relative extinction).

23 The MCMC sampler relies on emcee ([Foreman-Mackey et al., 2013, 2019](#)), while two options
24 are available for nested sampling: nestle ([Barbary, 2013](#)) and ultranest ([Buchner, 2021b](#)).
25 The samplers have been adapted for flexibility - they are usable on any grid of input models
26 provided by the user, simply requiring a snippet function specifying the format of the input.
27 Moreover they can sample the effect of blackbody component(s) (either as a separate model or
28 as extra components to an atmospheric model), extinction, and different extinction laws than
29 ISM. The samplers can accept either uniform or Gaussian priors for each model parameter.
30 In the case of the MCMC sampler, a prior on the mass of the object can also be provided
31 if surface gravity is one of the model parameters. The code also considers convolution and
32 resampling of model spectra to match the observed spectrum. Either spectral resolution or
33 photometric filter transmission (or combinations thereof for compound input spectra) can be
34 provided as input to the algorithm, for appropriate convolution/resampling of different parts of
35 the model spectrum. The adopted log-likelihood expression can include i) spectral covariance
36 between measurements of adjacent channels of a given instrument, and ii) additional weights
37 between measurements of adjacent channels of a given instrument, and ii) additional weights
38 between measurements of adjacent channels of a given instrument, and ii) additional weights

³⁹ that are proportional to the relative spectral bandwidth of each measurement, in case these
⁴⁰ are obtained from different instruments (e.g. photometry+spectroscopy):

$$\log \mathcal{L}(D|M) = -\frac{1}{2} [\mathbf{W}(\mathbf{F}_{\text{obs}} - \mathbf{F}_{\text{mod}})^T] \mathbf{C}^{-1} [\mathbf{W}^T(\mathbf{F}_{\text{obs}} - \mathbf{F}_{\text{mod}})] \quad (1)$$

⁴¹ where \mathbf{F}_{obs} and \mathbf{F}_{mod} are the fluxes of the observed and model spectra respectively, \mathbf{C} is the
⁴² spectral covariance matrix, and \mathbf{W} is the vector of weights $w_i \propto \Delta\lambda_i/\lambda_i$, with $\Delta\lambda_i$ the width
⁴³ of spectral channels (for integral field spectrograph points) or the FWHM of photometric
⁴⁴ filters.

⁴⁵ A jupyter notebook tutorial illustrates most available features in `special` through their
⁴⁶ application for the analysis of the composite spectrum of CrA-9 B/b (Christiaens et al., 2021).
⁴⁷ It is available on [GitHub](#), [Binder](#) and the [documentation](#) of `special`.

⁴⁸ Acknowledgements

⁴⁹ VC acknowledges financial support from the Belgian F.R.S.-FNRS.

⁵⁰ References

- ⁵¹ Allers, K. N., Jaffe, D. T., Luhman, K. L., Liu, M. C., Wilson, J. C., Skrutskie, M. F., Nelson, M., Peterson, D. E., Smith, J. D., & Cushing, M. C. (2007). Characterizing Young Brown Dwarfs Using Low-Resolution Near-Infrared Spectra. *657*(1), 511–520. <https://doi.org/10.1086/510845>
- ⁵⁵ Barbary, K. (2013). `nestle`. In *Github repository*. GitHub. <https://github.com/kbarbary/nestle>
- ⁵⁷ Buchner, J. (2021a). Nested Sampling Methods. *arXiv e-Prints*, arXiv:2101.09675. <https://arxiv.org/abs/2101.09675>
- ⁵⁹ Buchner, J. (2021b). UltraNest - a robust, general purpose Bayesian inference engine. *The Journal of Open Source Software*, 6(60), 3001. <https://doi.org/10.21105/joss.03001>
- ⁶¹ Christiaens, V., Ubeira-Gabellini, M.-G., Cánovas, H., Delorme, P., Pairet, B., Absil, O., Casassus, S., Girard, J. H., Zurlo, A., Aoyama, Y., Marleau, G.-D., Spina, L., van der Marel, N., Cieza, L., Lodato, G., Pérez, S., Pinte, C., Price, D. J., & Reggiani, M. (2021). A faint companion around CrA-9: protoplanet or obscured binary? *MNRAS*, 502(4), 6117–6139. <https://doi.org/10.1093/mnras/stab480>
- ⁶⁶ Delorme, P., Schmidt, T., Bonnefoy, M., Desidera, S., Ginski, C., Charnay, B., Lazzoni, C., Christiaens, V., Messina, S., D’Orazi, V., Milli, J., Schlieder, J. E., Gratton, R., Rodet, L., Lagrange, A.-M., Absil, O., Vigan, A., Galicher, R., Hagelberg, J., ... Wildi, F. (2017). In-depth study of moderately young but extremely red, very dusty substellar companion HD 206893B. *608*, A79. <https://doi.org/10.1051/0004-6361/201731145>
- ⁷¹ Feroz, F., Hobson, M. P., & Bridges, M. (2009). MULTINEST: an efficient and robust Bayesian inference tool for cosmology and particle physics. *398*(4), 1601–1614. <https://doi.org/10.1111/j.1365-2966.2009.14548.x>
- ⁷⁴ Foreman-Mackey, D., Farr, W., Sinha, M., Archibald, A., Hogg, D., Sanders, J., Zuntz, J., Williams, P., Nelson, A., de Val-Borro, M., Erhardt, T., Pashchenko, I., & Pla, O. (2019). emcee v3: A Python ensemble sampling toolkit for affine-invariant MCMC. *The Journal of Open Source Software*, 4(43), 1864. <https://doi.org/10.21105/joss.01864>
- ⁷⁸ Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. (2013). emcee: The MCMC Hammer. *125*(925), 306. <https://doi.org/10.1086/670067>

- 80 Goodman, J., & Weare, J. (2010). Ensemble samplers with affine invariance. *Communications
81 in Applied Mathematics and Computational Science*, 5(1), 65–80. [https://doi.org/10.
82 2140/camcos.2010.5.65](https://doi.org/10.2140/camcos.2010.5.65)
- 83 Gorlova, N. I., Meyer, M. R., Rieke, G. H., & Liebert, J. (2003). Gravity Indicators in the
84 Near-Infrared Spectra of Brown Dwarfs. 593, 1074–1092. <https://doi.org/10.1086/376730>
- 85 Greco, J. P., & Brandt, T. D. (2016). The Measurement, Treatment, and Impact of Spectral
86 Covariance and Bayesian Priors in Integral-field Spectroscopy of Exoplanets. 833, 134.
87 <https://doi.org/10.3847/1538-4357/833/2/134>
- 88 Mukherjee, P., Parkinson, D., & Liddle, A. R. (2006). A Nested Sampling Algorithm for
89 Cosmological Model Selection. 638(2), L51–L54. <https://doi.org/10.1086/501068>
- 90 Skilling, J. (2004). Nested Sampling. In R. Fischer, R. Preuss, & U. V. Toussaint (Eds.),
91 *Bayesian inference and maximum entropy methods in science and engineering: 24th inter-
92 national workshop on bayesian inference and maximum entropy methods in science and
93 engineering* (Vol. 735, pp. 395–405). <https://doi.org/10.1063/1.1835238>
- 94 Slesnick, C. L., Hillenbrand, L. A., & Carpenter, J. M. (2004). The Spectroscopically
95 Determined Substellar Mass Function of the Orion Nebula Cluster. 610, 1045–1063.
96 <https://doi.org/10.1086/421898>

DRAFT